



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



General review

Migraine treatment: Position paper of the French Headache Society

X. Moisset^{a,1,*}, G. Demarquay^{b,c,1}, S. de Gaalon^d, C. Roos^e, A. Donnet^f,
 P. Giraud^g, E. Guégan-Massardier^h, C. Lucasⁱ, J. Mawet^e, D. Valade^j,
 V. Corand^k, C. Gollion^l, N. Moreau^{m,n}, L. Grangeon^h,
 M. Lantéri-Minet^{a,o}, A. Ducros^p

^a Université Clermont Auvergne, CHU Clermont-Ferrand, Inserm, Neuro-Dol, 63000 Clermont-Ferrand, France

^b Hospices Civils de Lyon, University of Lyon, Lyon, France

^c Lyon Neuroscience Research Center (CRNL), Inserm U1028, CNRS UMR5292, Lyon, France

^d Department of Neurology, hôpital Laënnec, CHU de Nantes, Nantes, France

^e Emergency Headache Center (Centre d'Urgences Céphalées), Department of Neurology, Lariboisière Hospital, Assistance publique des Hôpitaux de Paris, Paris, France

^f Pain assessment and treatment centre, FHU INOVRAIN, hôpital de La Timone, Marseille, France

^g Department of Neurology, Annecy Genevois Hospital, Annecy, France

^h Department of Neurology, Rouen University Hospital, Rouen, France

ⁱ Pain assessment and treatment centre, Service de Neurochirurgie, Hôpital Salengro, CHU de Lille, Lille, France

^j Department of Neurosurgery, hôpital Pitié-Sapêtrière, Paris, France

^k Pain consultation, Polyclinique Jean-Villar, 33520 Bruges, France

^l Neurology Department, CHU de Toulouse, Toulouse, France

^m Orofacial neurobiology laboratory, EA 7543, Université Paris Cité, Paris, France

ⁿ Hôpital Bretonneau, Service de médecine buccodentaire, AP-HP, Paris, France

^o Pain Department and FHU InoPain, CHU de Nice, Côte Azur Université, Nice, France

^p Department of Neurology, Gui-de-Chauliac Hospital, CHU de Montpellier, University of Montpellier, 34000 Montpellier, France

INFO ARTICLE

Keywords:

Migraine

Headache

Guidelines

Recommendations

CGRP

Gepants

ABSTRACT

The French migraine management recommendations were published in 2021. However, in the last three years, new data have come to light and new drugs have been approved (eptinezumab, rimegepant and atogepant) by the European Medicines Agency that require us to take a position on their use and to update certain elements of the recommendations. The first important message concerns the position of the French Headache Society on the use of preventive treatments (monoclonal antibodies and gepants) targeting the calcitonin gene-related peptide (CGRP) pathway. In terms of efficacy and safety, and as suggested by other national headache societies, these treatments can be offered as first-line treatment, although the scope defined by the French national health authority for possible reimbursement is limited to patients with severe migraine, at least eight headache days per month and

* Corresponding author. Service de Neurologie, CHU Gabriel-Montpied, 58, rue Montalembert, 63000 Clermont-Ferrand, France.

E-mail address: xavier.moisset@gmail.com (X. Moisset).

¹ Contributed equally.

<https://doi.org/10.1016/j.neurol.2024.09.008>

0035-3787/© 2024 The Author(s). Published by Elsevier Masson SAS. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

for whom two previous preventive treatments have failed. Another important change concerns the position of topiramate as a preventive treatment for migraine in women of childbearing age. This treatment has been proposed as a first-line treatment for chronic migraine. However, recent pharmacovigilance data have highlighted a potential adverse effect on neurodevelopment in children exposed in utero. As a result, this treatment is formally contraindicated during pregnancy and must be used with extreme caution in women of childbearing age (effective contraception, no therapeutic alternative available and annual follow-up as with valproate). It can therefore no longer be offered as first-line treatment for women of childbearing age.

© 2024 The Author(s). Published by Elsevier Masson SAS. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Since the publication of the latest French recommendations for the management of migraine in adults, concerning diagnosis [1], drug treatments [2] and non-drug therapies [3], a number of important points need to be clarified. It is important to underline that the management of migraine has evolved considerably in recent years, thanks to improved knowledge of the pathophysiology of migraine. Indeed, the discovery in the 1980s of the key role played by calcitonin gene-related peptide (CGRP) in the development of migraine pain led to the development of new targeted therapies [4,5]: monoclonal antibodies targeting CGRP or its receptor, and non-peptide small molecule CGRP receptor antagonists known as gepants. The main point of this article refers to the position taken by the French Headache Society on the use of the four monoclonal antibodies targeting the CGRP pathway (in alphabetical order, eptinezumab, erenumab, fremanezumab, galcanezumab) and the gepants. With the arrival of atogepant and rimegepant on the French market, these compounds must be positioned in the therapeutic armamentarium. Another very important point concerns the potential neurodevelopmental risk highlighted for topiramate, which was not clearly known when the 2021 guidelines were published and which now justifies extreme caution when using this drug in women of childbearing age. Finally, the marketing discontinuation of flunarizine is specified.

1. New data on monoclonal antibodies targeting the CGRP pathway and the position of the French headache society on their use and reimbursement

1.1. Update on the place of the four monoclonal antibodies

As for monoclonal antibodies targeting the CGRP pathway, only three (erenumab, fremanezumab and galcanezumab) were approved by the European Medicines Agency (EMA) for migraine prophylaxis at the time of publication of the 2021 recommendations. Today, eptinezumab is also approved by the EMA with the same indication as the other three, namely prevention of migraine in adults with at least four migraine days per month. Unfortunately for patients, none of these four treatments are reimbursed at the time of writing.

Up to now, fremanezumab and galcanezumab are available in pharmacies. Only eptinezumab, given intravenously every three months, is currently administered in some French hospitals and clinics, generally free of charge for the patient, with a specific status known in French as “reserve hospitalière”. Each hospital has the choice of whether or not to list new treatments, and decisions are not uniform across all French healthcare establishments. What’s more, for hospitals that do provide access to this treatment, only the dose of 100 mg per infusion is available, making it impossible to increase the dosage to 300 mg per infusion for non-responders. Based solely on the efficacy and safety data for antibodies targeting the CGRP pathway, these four treatments can be offered as first-line treatment to all patients presenting with at least four migraine days per month, according to their approval and as proposed by the European Headache Federation [6] and the American Headache Society [7]. In line with EMA decision, prescription is authorized in France under the terms of their approval, i.e. for all patients with at least four migraine days per month. However, the scope defined by the French national health authority (HAS standing for Haute Autorité de santé) for possible reimbursement is limited to patients with severe migraine, at least eight headache days per month and for whom two previous preventive treatments have failed. This rule should therefore be respected for patients receiving infusions of eptinezumab with the cost at least partly covered by the hospital supplying the product, but does not apply to patients who are prepared to finance the whole cost of treatment themselves, for galcanezumab and fremanezumab. These rules will also have to be followed for subcutaneous antibodies if they are reimbursed in the coming years. From a medico-economic point of view, a study carried out in the United Kingdom showed that eptinezumab is a cost-effective treatment in patients with four or more migraine days per month and for whom three or more other preventive migraine treatments have failed [8]. Erenumab is also cost-effective for episodic [9] and for chronic [10] migraine.

It is important to note that the notion of severe migraine, taken into account in the HAS wording, was proposed in 2021 [11] but that modifications were suggested in 2023 after the criteria were tested in the general patient population (Table 1) [12]. The two new proposals must necessarily be tested in turn in order to validate the best definition.

Table 1 – Definitions of severe migraine.

Initial definition proposed in 2021 [11]	Headaches fulfilling either criterion A or criterion B for at least three months A. Headache frequency of at least eight migraine days per month B. Headache frequency < 8 migraine days per month, but associated with at least one of the following criteria: 1. Headache impact test-6 (HIT-6) score ≥ 60 2. Necessitating complete interruption of activity for 50% of headaches
Modified definition proposed in 2023 [12]	Headaches fulfilling either criterion A or criterion B for at least three months A. Headache frequency of at least eight migraine days per month B. Headache frequency < 8 migraine days per month, but associated with both of the following criteria: 1. HIT-6 score ≥ 60 2. Necessitating complete interruption of activity for 50% of headaches
Proposal 1	
Modified definition proposed in 2023 [12]	Headaches fulfilling either criterion A or criterion B for at least three months A. Headache frequency of at least eight migraine days per month B. Headache frequency < 8 migraine days per month but associated with a HIT-6 score ≥ 65
Proposal 2	

As with any patient for whom a preventive treatment is prescribed, a comprehensive management program should be applied to assess the frequency of migraine days (using an agenda), the impact of migraine on quality of life (e.g. using the HIT-6 scale), and any comorbid anxiety or depression (e.g. using the HAD questionnaire).

1.2. Update on efficacy

Up to now, there have been no randomized controlled trials directly comparing the four monoclonal antibodies. Indirect comparisons did not reveal any significant difference in efficacy between these four compounds [13]. However, it is important to note that response (or non-response) to one antibody is not predictive of response to another. Indeed, it has been shown that the switch from erenumab to another CGRP-mAb led to a $\geq 30\%$ response in more than one-third of the patients [14,15]. In addition, half of the non-responders to anti-CGRP mAbs at 12 weeks may be late responders according to an open-label study, which would justify, at least in some cases, evaluating efficacy at 24 weeks and extending therapy beyond 12 months [16]. Real-world data provides a better understanding of the proportion of patients who continue to benefit from treatment after it has been stopped [17].

Studies directly comparing anti-CGRP monoclonal antibodies with conventional treatments are scarce. One study demonstrated that erenumab was better tolerated than topiramate. A secondary endpoint compared the efficacy of the two treatments [18]. It clearly showed that erenumab was superior for both tolerance and efficacy. When it comes to indirect comparisons, they all point to a much better tolerability of this new therapeutic class compared with conventional treatments, particularly topiramate [13]. In terms of efficacy, the data are less clear, with some meta-analyses showing that topiramate is at least as effective as monoclonal antibodies [19], while others support the superiority of antibodies [13]. In addition, it has recently been shown that using CGRP mAbs earlier in the treatment of migraine patients is more effective than offering another oral medication [20].

Regarding factors predicting response to monoclonal antibodies targeting the CGRP pathway, a meta-analysis showed that a good response to triptans and unilateral pain with or without unilateral autonomic symptoms are predictors of a good response to these drugs [21]. Conversely, obesity,

interictal allodynia, higher migraine frequency, greater disability at baseline, a higher number of unsuccessful previous prophylactic medications and psychiatric comorbidities including depression are predictive of a poor response to monoclonal antibodies targeting the CGRP pathway [21,22]. It is important to emphasise that poor response does not mean absence of response. Treatments targeting the CGRP pathway should be considered in patients who may benefit from treatment despite predictors of poor response. For example, in a large French series of patients who had failed all previously available treatments (an average of 11 treatments tried), the introduction of erenumab treatment resulted in a 50% response in just over one in two patients [23].

To date, there is insufficient evidence to support or reject the efficacy of combining treatments targeting the CGRP pathway with oral migraine prophylactic agents [24]. Some real-life data are beginning to emerge concerning the combined use of an anti-CGRP antibody and botulinum toxin type A [25]. There seems to be a benefit to this type of combination, provided access to these therapies is available.

1.3. Update on treatment duration and tolerance

A large proportion of patients experience a significant worsening of their migraine frequency after stopping treatment, justifying resumption of treatment [26–28]. It therefore seems important to consider the possibility of long-term treatment for the majority of patients. Long-term efficacy of anti-CGRP antibodies is as good as short-term efficacy [29]. In the context of long-term treatment, the safety profile of these treatments still appears to be very good. However, there is a risk of arterial hypertension, which should be monitored in all patients [30,31]. A risk of Raynaud’s phenomenon has been identified [32] and a rare risk of alopecia has also been described [33]. A number of other potential adverse effects have been reported, including sexual dysfunction in both men [34] and women [35] and impaired wound healing [36]. It is important that clinicians pay close attention to these potential side effects to confirm whether they exist, even if they are very rare. On the other hand, the initial data available on exposure during pregnancy is reassuring, although the data is limited and does not yet allow us to recommend the use of these antibodies during pregnancy [37]. The risk of constipation and allergy is confirmed [38].

1.4. Position of the French Headache Society on reimbursement

Despite the safety and efficacy data presented above, and despite European marketing authorization, at the time of writing this article, these drugs are not reimbursed in France. Fig. 1 provides a simplified overview of the main elements involved in making a medicine available and reimbursed on the French market. For the four monoclonal antibodies (Table 2), the HAS concluded that the medical benefit (SMR for *service médical rendu*) is important in the preventive treatment of patients with severe migraine with at least eight migraine days per month, for whom at least two prophylactic treatments have failed and who have no cardiovascular problems (patients who have had a myocardial infarction, stroke, transient ischemic attack, unstable angina or coronary artery bypass surgery). At the same time, the HAS Transparency Commission concluded that there was no improvement in the medical benefit (ASMR V for *absence d'amélioration du service médical rendu*) compared with existing treatments. This decision was made in the absence of a direct comparative trial against a reference treatment (beta-blocker or topiramate or candesartan or amitriptyline). Reimbursement of the four monoclonal antibodies is therefore possible, provided that treatment costs are reduced. With reference treatments such as topiramate or beta-blockers costing less than €10 a month, it would seem difficult to offer a monoclonal antibody at a lower cost. On the other hand, the Commission's conclusion that these molecules offer no therapeutic advances is highly questionable. Indeed, all these molecules have shown efficacy even in patients for whom two to four prophylactic treatments had failed. Moreover, it is not technically feasible to conduct a therapeutic trial against a reference treatment in patients who have had two to four failed reference prophylactic treatments. The data available (in randomized clinical trials and in real-world studies) for the four antibodies in these populations

therefore show that these treatments represent a breakthrough in these difficult-to-treat patients [39–42].

In conclusion, the four monoclonal antibodies targeting the CGRP pathways are effective and well tolerated and should be offered to patients who need them. Taking into account only the EMA approval and the efficacy and safety data, it should be possible to offer these treatments targeting the CGRP pathways as first-line treatment for all patients who need preventive therapy. For economic reasons, the French Headache Society is in favor of reimbursing these treatments for patients with severe migraine if at least two of the other reference treatments (beta-blockers, candesartan, amitriptyline, topiramate) are ineffective, intolerable or contraindicated.

2. Arrival of gepants on the French market

Rimegepant has proven its effectiveness in the treatment of migraine attacks [43]. It has also proved useful as a preventive treatment [44]. It has been approved by the EMA as both an attack treatment and a preventive treatment for patients with episodic migraine with at least four migraine attacks per month, and can therefore be prescribed for both indications. It has been available on the French market since October 2023, but is not reimbursed. The pharmaceutical company marketing rimegepant did not consult the Transparency Commission and chose to market this treatment without claiming reimbursement. It therefore has no SMR or ASMR at the time of writing this article.

Atogepant, has demonstrated its efficacy in the preventive treatment of episodic and chronic migraine [45,46]. It has also been shown to be effective in patients with episodic migraine for whom two to four classes of conventional oral preventive treatment have failed [47]. This treatment was approved in August 2023 by the EMA to prevent migraines in adults who




Fig. 1 – Simplified representation of the factors taken into account for a drug to be eligible for reimbursement and for a reimbursement price to be defined.

Table 2 – Opinions from the European and French authorities on the four monoclonal antibodies targeting the CGRP pathway and two gepants, and their availability on the French market.

	EMA approval	CT opinion	Approval for communities ^b	CEPS opinion	Availability on the French market
Eptinezumab (anti-CGRP)	MA 24/01/2022, Migraine prevention in adults who have migraines at least 4 days a month	05/10/2022 SMR important ASMR V ^a	Yes	Price negotiations fail, no reimbursement for patients	For hospital use only, available in certain healthcare facilities at no cost to the patient
Erenumab (anti-CGRP receptor)	MA 26/07/2018 Migraine prevention in adults who have migraines at least 4 days a month	27/02/2019 revised on 30/03/2022 SMR important ASMR V ^a	Yes		Not available
Fremanezumab (anti-CGRP)	MA 28/03/2019 Migraine prevention in adults who have migraines at least 4 days a month	16/09/2020 revised on 14/09/2022 SMR important ASMR V ^a	Yes		Available in pharmacies, not reimbursed, recommended retail price €270
Galcanezumab (anti-CGRP)	MA 14/11/2018 Migraine prevention in adults who have migraines at least 4 days a month	24/06/2020 SMR important ASMR V ^a	Yes		Available in pharmacies, not reimbursed, recommended retail price €245
Rimegepant	MA 24/02/2022 Acute treatment of migraine with or without aura in adults Migraine prevention in adults who have migraines at least 4 days a month	Not requested by the company	No	Not requested by the company	Available in pharmacies, not reimbursed; €60–80 for 2 tablets
Atogepant	MA 25/08/2023 Migraine prevention in adults who have migraines at least 4 days a month	06/12/2023 SMR important ASMR V ^a	Yes	Discussion in progress	Available in pharmacies, not reimbursed; €250–300 for 28 tablets

EMA: European Medicines Agency; CT: commission de la transparence of the French health authority (HAS); CEPS: Comité Économique des Produits de Santé (Economic Committee for Health Products); MA: marketing authorization.

^a CT opinion for the four monoclonal antibodies: favourable opinion for reimbursement only as preventive treatment for migraine in patients with severe migraine with at least 8 migraine days per month, who have failed at least two preventive treatments and who have no cardiovascular problems.

^b In order to be purchased and used in hospitals, medicines must be on the list of speciality medicines approved for use by local authorities and various public services.

have migraines at least four days a month. In December 2023, the HAS has come out in favour of possible reimbursement only for patients with at least eight migraine days per month and at least two failed prophylactic treatments (SMR important, ASMR V). It has been available in France since June 2024. Based solely on the efficacy and safety, atogepant can be offered as first-line treatment to all patients presenting with at least four migraine days per month, but the administrative rules in France for possible reimbursement are the same as those that apply to antibodies targeting the CGRP pathway.

Other gepants, including ubrogepant and zavegepant nasal spray, have proven effective as attack treatments [48,49]. These gepants were approved by the United States Food and Drug Administration (FDA) but not yet by the EMA. A recent study showed that taking ubrogepant very early on, from the prodrome phase, was effective in treating the attack [50]. What's more, the current data seem to be very favorable for

gepants in terms of the risk of developing medication overuse headaches if taken repeatedly. That was shown in animals for olcegepant [51]. There are also human data with rimegepant and atogepant, which have been shown to be effective as prophylactic treatments and not to cause medication overuse headache. The half-life of these two drugs is around 11 hours. In the case of ubrogepant and zavegepant, the half-life is shorter (around six hours), which could theoretically modify the risk of headaches from overuse. However, the preclinical data for ubrogepant are entirely reassuring [52]. The gepants listed in this last paragraph are not currently available in France.

In conclusion, rimegepant (episodic migraine only) and atogepant are effective and well tolerated and should be offered to patients who need them. Taking into account only the EMA approval and the efficacy and safety data, it should be possible to offer these gepants as first-line treatment for all

patients who need preventive therapy (adults who have migraines at least four days a month) and rimegepant as an attack treatment. For economic reasons, the French Headache Society is in favor of reimbursing these treatments for patients with severe migraine if at least two of the other reference preventive treatments (beta-blockers, candesartan, amitriptyline, topiramate) are ineffective, intolerable or contraindicated. The French Headache Society also supports the reimbursement of rimegepant for the treatment of attacks in patients who have received failed regimens, have a contraindication to, or are intolerant to both nonsteroidal anti-inflammatory drugs (NSAIDs) and triptans.

3. Update on the management of non-responders to triptans

Non-response to triptan has been defined recently and proposed criteria are presented in Table 3 [53]. However, it has been shown that some patients who do not respond to two triptans may respond well to a third or even a fourth triptan [54]. This study, using the German registry, showed that the highest response rates were seen with nasal and oral zolmitriptan, oral eletriptan and subcutaneous sumatriptan. Acute treatment optimization might include switching to one of the triptans with the highest response rates. Attack treatments are listed in Tables 4A and 4B with their usual doses, the main adverse effects and the main contraindications. Prophylactic treatments are listed in Tables 5A and 5B.

As proposed in the 2021 recommendations, another option for optimising attack treatment is to combine an NSAID and a triptan, if necessary with an anti-emetic (metoclopramide in particular). The use of a fixed combination of sumatriptan and naproxen has been shown to be superior to each of the compounds used in isolation, and deserves to be tried (not commercially available in France at time of writing) [55]. In fact, this fixed combination is the only one for which superiority over NSAIDs and triptan taken alone has been demonstrated in large randomized controlled trials. However, other combinations of NSAIDs and triptan may be of interest, and different combinations may be tried.

Another solution for optimizing attack treatment could be to change the therapeutic class and use a gepant. Rimegepant should be of interest to patients who do not respond optimally to triptans, or who have an adverse reaction or contraindication to both NSAIDs and triptans, especially as tolerance is excellent [56]. However, no studies performed specically on triptans/NSAIDs non-responders have been published. The

results of the trial (NCT05509400) are eagerly awaited to see if they confirm this expectation.

Although there are no clearly defined criteria for declaring a patient resistant or refractory to an attack treatment with NSAIDs or gepant, it seems legitimate to try several molecules in the same class to optimize attack treatment.

4. New data on the risk of certain pharmacological treatments during pregnancy

Topiramate is an effective treatment of episodic and chronic migraine. It is nevertheless important to note that recent meta-analyses clearly show that the effect in chronic migraine is less clear-cut than in episodic migraine and that treatments with monoclonal antibodies targeting the CGRP pathway or with botulinum toxin type A are more effective than topiramate in chronic migraine [19,57]. Real-world data on the use of botulinum toxin type A show good efficacy and safety [58]. Botulinum toxin type A can be used as a preventive treatment for chronic migraine, even in patients with a vascular contraindication to the use of CGRP-targeting treatments. There is insufficient data to support the use of botulinum toxin during pregnancy, although the data on exposure during the first trimester appear reassuring [59]. With regard to topiramate, recent data suggest that its use in pregnant women carries a risk of neurodevelopmental problems in the unborn child [60]. According to a large and well-conducted study, there is an increased risk of autism spectrum disorder, with a hazard ratio of 2.8 (95% CI: 1.4–5.7), as well as a risk of intellectual disability with a hazard ratio [95% CI] of 3.5 [1.4–8.6]. The effect size for this risk appears to be at least equivalent to that of valproate, whose risk is already well known. Consequently, the risks of neurodevelopmental disorders for an unborn child must be clearly explained to women of childbearing age. Following these data, the French national drug safety agency (Agence nationale de sécurité du médicament [ANSM]) has modified the conditions for prescribing and dispensing topiramate to girls, adolescents and women of childbearing age, with treatment initiation reserved for neurologists and an annual care agreement co-signed by the patient and the neurologist [61]. We therefore no longer recommend topiramate as first-line treatment in migraine for women of childbearing age. The same conditions have applied to valproate prescriptions since 2017. Thus, valproate is also not recommended for women of childbearing age if other therapeutic options are available.

It is important to note that all the data for valproate are consistent, with a constant excess risk in all the latest major studies, whether it is the risk of autism, intellectual disability or the development of psychiatric disorders [62,63]. It is not the case for topiramate. Indeed, the most recent study [62], which only looked at the risk of autism, found no association between in utero exposure to topiramate and the risk of developing an autism spectrum disorder at the age of 8. In fact, the association was substantially attenuated for topiramate after adjustment for indication and other confounders.

As a reminder, beta-blockers and amitriptyline can be used during pregnancy, although migraine attacks are usually rare. Both treatments have a high level of efficacy for episodic

Table 3 – Definition of resistant and refractory patients for triptans.	
Triptans [53]	Responder when one or several triptans lead to effective acute treatment in at least three out of four migraine attacks
	Resistant in the presence of failure of at least 2 triptans
	Refractory in the presence of failure of at least 3 triptans, including subcutaneous sumatriptan 6 mg/0.5 mL

Table 4A – Nonspecific acute migraine treatments.

	Dose, route	Main side effects	Main contraindications ^a
Analgesics (French Market Approval for the acute treatment of migraine, yes or no)			
Paracetamol (no)	500, 1000 mg (tablet) Maximum 4 g/day	Paracetamol: hepatic and hematologic toxicity	Severe hepatic insufficiency
Paracetamol + caffeine (no)	500 mg + 50 mg (tablet) Maximum 6 tablets/day	Caffeine: palpitation insomnia	
NSAIDs			
Diclofenac (no)	25, 50, 100 mg (tablet) Maximum 150 mg/day	Hemorrhagic syndrome Digestive disorder, dyspepsia, nausea, diarrhea, constipation Dizziness, asthenia	Active gastroduodenal ulcer Hypersensitivity to NSAIDs Hemorrhagic risk (cerebral, digestive other), Severe hepatic or renal insufficiency Pregnancy (after the 5th month)
Flurbiprofen (no)	8.75 mg (tablet) Maximum 5 tablets/day		
Ibuprofen (yes)	200, 400 mg (tablet) Maximum 1200 mg/day		
Indomethacin (no)	25, 75 mg (tablet) 100 mg (suppository) Maximum 300 mg/day		
Ketoprofen (yes)	100, 150 mg (tablet) 100 mg (suppository) Maximum 200 mg/day		
Naproxen (no)	550, 1000 mg (tablet) Maximum 1100 mg/day		
Acetylsalicylate acid, aspirin (no)	1000 mg (tablet, powder, disintegrating tablet) Maximum 3000 mg/day	Acetylsalicylate: digestive disorder, hemorrhage, allergy, Reye syndrome	Acetylsalicylate: active gastroduodenal ulcer, hemorrhagic risk (cerebral, digestive other), pregnancy, asthma, severe hepatic, cardiac or renal insufficiency, hypersensitivity, pregnancy
Acetylsalicylate + metoclopramide (yes)	900 mg + 10 mg (powder) Maximum 3/day	Metoclopramide: dyskinetic syndrome, restlessness psychiatric disorder, endocrine disorder	Metoclopramide: gastrointestinal hemorrhage, digestive perforation, history of dyskinesia, extrapyramidal syndrome, children
^a The contraindications and adverse reactions are not exhaustive, but are listed in order of frequency of occurrence. Interactions are not listed. See Vidal.			

migraine [2]. In chronic migraine, the efficacy of propranolol has been considered fair [2], but a recent study showed that propranolol is as effective as topiramate in this indication [64]. Propranolol may therefore be a reasonable first-line option for patients with chronic migraine.

An alert published in August 2023 by the ANSM highlighted the potential risk for a child born to a father who had been exposed to valproate. In 2018, the EMA asked laboratories marketing valproate or its derivatives to conduct studies to better clarify the risks of these treatments. The results of one

such study (EUPAS34201, presented in May 2023) suggested an increased risk of neurodevelopmental disorders in children whose epileptic fathers had been treated in the three months prior to conception with valproate or its derivatives. This risk ranged from 5.6% to 6.3% in children whose fathers had been exposed to valproate, versus 2.5% to 3.6% in those born to fathers treated with lamotrigine or levetiracetam [65]. It is therefore recommended from a legal point of view to inform men of this potential risk during the three months (spermatogenesis lasting an average of nine weeks) preceding

Table 4B – Specific acute migraine treatments.

	Dose (route)	Main side effects	Main contraindications ^a
Triptans (French Market Approval for the acute treatment of migraine, yes or no)			
Almotriptan (yes)	12.5 mg (tablet) Maximum 25 mg/day	Paresthesia of extremities, nausea, feeling of cold, dizziness, asthenia, “chest syndrome” (feeling of constriction in the chest and neck), flushing, somnolence Rare cases of coronary spasms, severe hypertension, serotonin syndrome	Coronary heart disease Wolff Parkinson White syndrome Myocardial infarction Peripheral arterial disease Raynaud TIA and stroke Uncontrolled hypertension Serious hepatic or renal insufficiency Concurrent treatment with a MAO inhibitor Cross allergy with sulfonamides (except for rizatriptan and zolmitriptan)
Eletriptan (yes)	20 or 40 mg (tablet) Maximum 80 mg/day		
Frovatriptan (yes)	2.5 mg (tablet) Maximum 5 mg/day		
Naratriptan (yes)	2.5 mg (tablet) Maximum 5 mg/day		
Rizatriptan (yes)	5, 10 mg (tablets), 10 mg (disintegrating tablet) Maximum 20 mg/day		
Sumatriptan (yes)	50 mg (tablets) Maximum 300 mg/day 10/20 mg (nasal spray) Maximum 40 mg/day 6 mg (subcutaneous injection) Maximum 12 mg/ day		
Zolmitriptan (yes)	2.5 mg (tablet/ disintegrating tablet) Maximum 10 mg/day Nasal spray 5 mg (not available in France)		
Gepants			
Rimegepant (yes, June 2023)	75 mg (tablet) Maximum 75 mg/day	Nausea Rare severe allergic reaction	History of hypersensitivity reaction to rimegepant
Ubrogepant (yes, January 2024)	50 mg, 100 mg (tablets) Maximum 200 mg/day	Nausea, drowsiness Rare severe allergic reaction	History of hypersensitivity reaction to Ubrogepant Concomitant use with a potent inhibitor of cytochrome P450 3A4 (CYP3A4)
Zavegepant (not available in France in 2024)	10 mg (nasal spray) Maximum 10 mg/day	Unusual taste, nausea, nasal discomfort, vomiting Rare severe allergic reaction	History of hypersensitivity reaction to zavegepant Severe (Child-Pugh class C) hepatic impairment Renal impairment (< 30 mL/min)
Ditans			
Lasmiditan (not available in France in 2024)	50 mg, 100 mg (tablets) Maximum 200 mg/day No more than one dose should be taken in 24 hours (FDA)	Common (> 2%): dizziness, fatigue, paresthesia, sedation, nausea and/or vomiting, muscle weakness. Significant driving impairment Central nervous system depression (dizziness, sedation) Rare (1%): hallucinations, euphoria Risk of misuse or abuse Rare cases of serotonin syndrome	Should be used with caution if used in combination with alcohol, cannabis or other CNS depressants No driving within the first 8 hours after intake (FDA)

^a The contraindications and adverse reactions are not exhaustive, but are listed in order of frequency of occurrence. Interactions are not listed.

Table 5A – Oral prophylactic treatments: dosage, side effects and contraindications.

Treatment (French Market Approval, yes or no)	Daily dosage Minimum–maximum (mean daily dosage)	Main side effects	Main contraindications ^a
Amitriptyline (yes)	10–100 mg (25 mg) Once at dinner time	Dry mouth, somnia, weight gain	Absolute: glaucoma, prostatic adenoma Relative: obesity
Beta-blocker			
Propranolol (yes)	20–240 mg (80 mg) BID or once in the morning (extended release)	Common: asthenia, poor tolerance to effort Rare: depression	Absolute: asthma, heart failure, atrio- ventricular block, bradycardia Relative: depression
Metoprolol (yes)	50–200 mg (100 mg) Once in the morning (extended release)		
Nebivolol (no)	5–10 mg (10 mg) Once in the morning		
Atenolol (no)	50–200 mg (100 mg) Once in the morning		
Timolol (no)	10–60 mg (20 mg) BID		
Candesartan (no)	8–32 mg (16 mg) BID or once a day	Hypotension	Absolute: heart failure, renal artery stenosis, renal impairment, pregnancy Relative: hypotension
Gepant			
Atogepant (yes)	60 mg, once a day	Constipation, nausea	Severe (Child-Pugh class C) hepatic impairment
Rimegepant (yes)	75 mg every other day	Nausea Rare severe allergic reaction	History of hypersensitivity reaction to rimegepant
Lisinopril (no)	5–40 mg (20 mg) Once a day	Hypotension, dry cough, exanthema, impaired renal function	Angio-edema, renal artery stenosis, renal impairment, hyperkalemia, pregnancy
Lamotrigine (no)	25–300 mg (100 mg) Once or twice a day	Common: dizziness, insomnia Rare: serious hypersensitivity reactions, depression, suicidal ideation	Absolute: hypersensitivity to lamotrigine, breastfeeding Relative: previous allergy to another antiepileptic
Levetiracetam (no)	500–3000 mg Twice a day	Irritability, depression	Relative: renal impairment
Oxetorone (yes)	60–180 mg (120 mg) Once in the evening	Common: somnolence Rare: diarrhea, parkinsonism	Parkinson disease, parkinsonism, pregnancy
Pizotifene (yes)	50–300 mg (150 mg) BID	Common: sedation, weight gain	Obesity, glaucoma, prostatic adenoma, pregnancy
Topiramate (yes)	50–200 mg (100 mg) Once or twice a day	Common: paresthesia, weight loss, cognitive effects (word-finding difficulties), depression Rare: renal calculi, acute myopia with secondary angle closure glaucoma	Absolute: hypersensitivity to topiramate, pregnancy, glaucoma, severe pulmonary disease, metformin use, hepatic disease, nephrolithiasis, renal failure Do not use as first-line in women of childbearing age, annual care agreement if indication accepted Relative: depression, suicidal ideation
Valproate (no)	250–2000 mg (750 mg) Once in the evening or twice a day	Common: nausea, weight gain, somnolence, tremor, alopecia, augmentation des ASAT ALAT, hepatitis	Absolute: liver disease, pregnancy, mitochondrial disease Do not use in women of childbearing age, caution in male of childbearing potential (to be confirmed) Relative: obesity
Venlafaxine (no)	37.5–300 mg (75–150 mg) Once a day	Common: nausea, dry mouth, hyperhidrosis	Hypersensitivity to venlafaxine

^a The contraindications and adverse reactions are not exhaustive, but are listed in order of frequency of occurrence. Interactions are not listed.

Table 5B – Injectable prophylactic treatments: dosage, side effects and contraindications.

Active component (French Market Approval, yes or no)	Daily dosage Minimum–maximum (mean daily dosage)	Side effects	Contraindications ^a
OnabotulinumtoxinA (yes)	31–39 injections (155–195 UI) in 7 muscular groups, quarterly	Injection site pain	Absolute: myasthenia gravis, amyotrophic lateral sclerosis
Anti-CGRP or CGRP receptor antibodies Erenumab (yes)	70–140 mg SC monthly	Injection site pain or redness, constipation, increase in blood pressure, allergy, alopecia, Raynaud phenomenon	Myocardial infarction, stroke, TIA, uncontrolled vascular risk factor Pregnancy
Eptinezumab (yes)	100–300 mg IV quarterly		
Fremanezumab (yes)	225 mg SC monthly 675 mg SC quarterly		
Galcanezumab (yes)	240 mg SC the first month, then 120 mg SC monthly		

^a The contraindications and adverse reactions are not exhaustive, but are listed in order of frequency of occurrence. Interactions are not listed.

conception. However, the only study published so far is reassuring [66]. Further data is required to clarify this risk.

5. Marketing discontinuation of flunarizine from the French market

The Janssen-Cilag laboratory decided to stop marketing flunarizine at the end of October 2023. The withdrawal of this treatment from the market is therefore not a decision by the health authorities, but a commercial decision by the laboratory. This calcium antagonist also has anti-histaminic, anti-cholinergic and anti-dopaminergic effects. This treatment, which had proved its efficacy in several rather old randomized controlled trials [67], was approved in France as a second-line migraine treatment, providing relief for some patients. However, tolerance problems were not uncommon, and it was recommended not to use it for longer than six months. Thus, the level of recommendation was only moderate. Although other molecules are available for the preventive treatment of migraine, this withdrawal has a negative impact on the quality of life of some patients.

6. Conclusion

In conclusion, the four monoclonal antibodies targeting the CGRP pathway are effective and well tolerated and should be offered as preventative treatments to patients who need them. That is also the case for rimegepant (episodic migraine only) and atogepant. Taking into account only the EMA approval and the efficacy and safety data, it should be possible to offer these treatments targeting the CGRP pathways as first-line treatment for all patients who need preventive therapy. For economic reasons, the French Headache Society is in favor of reimbursing these treatments only for patients with severe migraine if at least two of the other reference preventive treatments (beta-blockers, candesartan, amitriptyline, topiramate) are ineffective, intolerable or contraindicated. So far, there is little evidence to rank CGRP mAbs, atogepant, and botulinum toxin in the

treatment of chronic migraine. Gepants are well tolerated and may be useful as an acute treatment. For economic reasons, the French Headache Society also supports the reimbursement of rimegepant for the treatment of attacks in patients who have received a failed regimen, have a contraindication to, or are intolerant to both NSAIDs and triptans. Topiramate should not be used as first-line treatment in women of childbearing age (annual care agreement to be co-signed by the patient). This is obviously the case also for valproate in women, but recent data also show a potential risk in children born to fathers exposed within three months of conception.

Disclosure of interest

X. Moisset has received financial support from Allergan-Abbvie, Amgen-Horizon, Aptyspharma, Biogen, BMS, Dr Reddy's, Grünenthal, Lilly, Lundbeck, Teva, Merck-Serono, Novartis, Orion, Pfizer, Roche, Sanofi-Genzyme and Sun Pharma and non-financial support from SOS Oxygène not related to the submitted work. X. Moisset is associate editor of *Pain Reports* and of *Revue Neurologique*.

G. Demarquay has received financial support from Allergan-Abbvie, Lilly, Lundbeck, Novartis, Pfizer Teva not related to the submitted work.

S. de Gaalon has received financial support from Allergan-Abbvie, Bayer, Boehringer, Lilly, Linde, Lundbeck, Novartis, Teva and BMS-Pfizer not related to the submitted work.

C. Roos has received financial support from Allergan-Abbvie, Biogen, Orkyn, Novartis, Mylan, Homeperf, Lilly, Teva, Ophtalmic, Lundbeck, Sanofi, Pfizer not related to the submitted work.

A. Donnet has received financial support from Abbvie/Allergan, Amgen, EliLilly, Lundbeck, Novartis, Perfood, Pfizer, and Teva not related to the submitted work.

P. Giraud has received financial support from Lundbeck, Merk, Pfizer and Sanofi.

E. Guégan-Massardier has received financial support from Allergan-Abbvie, Lundbeck, Novartis, Teva and BMS-Pfizer and Medtronic not related to the submitted work.

C. Lucas has received financial support from Abbvie, Homeperf, Grunenthal, Eli Lilly, Lundbeck, Orion, Pfizer, SOS Oxygène and Teva not related to the submitted work. C. Lucas is past-President of the SFEMC.

J. Mawet has received financial supports for boards, meeting presentations or travel expenses for congress from Lilly, Novartis, Abbvie, TEVA, Lundbeck, Pfizer, Orion Pharma, SOS Oxygène, Dr Reddy's and GEP Santé.

D. Valade and L. Grangeon declare that they have no competing interest.

V. Corand has received financial supports from Allergan-Abbvie, Lundbeck, Teva, BMS and Pfizer.

C. Gollion received direct honorarium for board membership and/or speak from Novartis, Teva, Lilly, Pfizer, Abbvie, Lundbeck and Orkyn, and grants from IHS and SFETD/APICIL for academic research, not related to the submitted work.

N. Moreau has received financial support from Grünenthal not related to the submitted work.

M. Lanteri-Minet has received financial support from Abbvie/Allergan, Amgen, Astellas, ATI, Boehringer, Boston Scientific, Glaxo-SmithKline, Grunenthal, Eli Lilly, Lundbeck, Medtronic, Menari, MSD, Novartis, Orion Pharma, Perfood, Pfizer, ReckittBenckiser, Saint-Jude, Sanofi-Aventis, Teva, UCB, UPSA and Zambon not related to the submitted work. M. Lanteri-Minet is associate editor of *Journal of Headache and Pain* and past-President of the SFEMC. The team of M. Lanteri-Minet in Nice University Hospital/Côte Azur University (FHU InovPain) received academic support from PHRCn and PHRCir and grants from Eli Lilly, Lundbeck, Novartis, Pfizer, Teva.

A. Ducros received direct honorarium for board membership and/or speaking from Novartis, Lundbeck, Teva, Pfizer, Abbvie, Lilly, SOS oxygène. A. Ducros is associate editor of *Cephalalgia*, and past-President of the SFEMC. The team of AD in Montpellier University received academic support from PHRCn and PHRCir, and grants from Pfizer for academic research programs.

REFERENCES

- [1] Demarquay G, Moisset X, Lanteri-Minet M, de Gaalon S, Donnet A, Giraud P, et al. Revised guidelines of the French Headache Society for the diagnosis and management of migraine in adults. Part 1: diagnosis and assessment. *Rev Neurol (Paris)* 2021;177:725–33. <http://dx.doi.org/10.1016/j.neurol.2021.07.001>.
- [2] Ducros A, de Gaalon S, Roos C, Donnet A, Giraud P, Guégan-Massardier E, et al. Revised guidelines of the French headache society for the diagnosis and management of migraine in adults. Part 2: pharmacological treatment. *Rev Neurol (Paris)* 2021;177:734–52. <http://dx.doi.org/10.1016/j.neurol.2021.07.006>.
- [3] Demarquay G, Mawet J, Guégan-Massardier E, de Gaalon S, Donnet A, Giraud P, et al. Revised guidelines of the French headache society for the diagnosis and management of migraine in adults. Part 3: non-pharmacological treatment. *Rev Neurol (Paris)* 2021;177:753–9. <http://dx.doi.org/10.1016/j.neurol.2021.07.009>.
- [4] Edvinsson L. CGRP and migraine: from bench to bedside. *Rev Neurol (Paris)* 2021;177:785–90. <http://dx.doi.org/10.1016/j.neurol.2021.06.003>.
- [5] Edvinsson L, Haanes KA, Warfvinge K, Krause DN. CGRP as the target of new migraine therapies – successful translation from bench to clinic. *Nat Rev Neurol* 2018;14:338–50. <http://dx.doi.org/10.1038/s41582-018-0003-1>.
- [6] Sacco S, Amin FM, Ashina M, Bendtsen L, Deligianni CI, Gil-Gouveia R, et al. European Headache Federation guideline on the use of monoclonal antibodies targeting the calcitonin gene related peptide pathway for migraine prevention – 2022 update. *J Headache Pain* 2022;23:67. <http://dx.doi.org/10.1186/s10194-022-01431-x>.
- [7] Charles AC, Digre KB, Goadsby PJ, Robbins MS, Hershey A, Society TAH. Calcitonin gene-related peptide-targeting therapies are a first-line option for the prevention of migraine: an American Headache Society position statement update. *Headache J Head Face Pain* 2024;64:333–41. <http://dx.doi.org/10.1111/head.14692>.
- [8] Griffin E, Shirley G, Lee XY, Awad SF, Tyagi A, Goadsby PJ. An economic evaluation of eptinezumab for the preventive treatment of migraine in the UK, with consideration for natural history and work productivity. *J Headache Pain* 2024;25:59. <http://dx.doi.org/10.1186/s10194-024-01749-8>.
- [9] Pozo-Rosich P, Poveda JL, Crespo C, Martínez M, Rodríguez JM, Irimia P. Is erenumab an efficient alternative for the prevention of episodic and chronic migraine in Spain? Results of a cost-effectiveness analysis. *J Headache Pain* 2024;25:40. <http://dx.doi.org/10.1186/s10194-024-01747-w>.
- [10] Khanal S, Underwood M, Naghdi S, Brown A, Duncan C, Matharu M, et al. A systematic review of economic evaluations of pharmacological treatments for adults with chronic migraine. *J Headache Pain* 2022;23:122. <http://dx.doi.org/10.1186/s10194-022-01492-y>.
- [11] Donnet A, Ducros A, Radat F, Allaf B, Chouette I, Lanteri-Minet M. Severe migraine and its control: a proposal for definitions and consequences for care. *Rev Neurol (Paris)* 2021;177:924–34. <http://dx.doi.org/10.1016/j.neurol.2020.11.012>.
- [12] Henri J, Redon S, Donnet A. Study of severe migraine in general medicine. *Rev Neurol (Paris)* 2023;179:993–9. <http://dx.doi.org/10.1016/j.neurol.2023.01.730>.
- [13] Lampl C, MaassenVanDenBrink A, Deligianni CI, Gil-Gouveia R, Jassal T, Sanchez-Del-Rio M, et al. The comparative effectiveness of migraine preventive drugs: a systematic review and network meta-analysis. *J Headache Pain* 2023;24:56. <http://dx.doi.org/10.1186/s10194-023-01594-1>.
- [14] Straube A, Broessner G, Gaul C, Hamann X, Hipp J, Kraya T, et al. Real-world effectiveness of fremanezumab in patients with migraine switching from another mAb targeting the CGRP pathway: a subgroup analysis of the Finesse Study. *J Headache Pain* 2023;24:59. <http://dx.doi.org/10.1186/s10194-023-01593-2>.
- [15] Overeem LH, Peikert A, Hofacker MD, Kamm K, Ruscheweyh R, Gendolla A, et al. Effect of antibody switch in non-responders to a CGRP receptor antibody treatment in migraine: a multi-center retrospective cohort study. *Cephalalgia* 2022;42:291–301. <http://dx.doi.org/10.1177/03331024211048765>.
- [16] Barbanti P, Aurilia C, Egeo G, Torelli P, Proietti S, Cevoli S, et al. Late response to anti-CGRP monoclonal antibodies in migraine: a multicenter prospective observational study. *Neurology* 2023;101:482–8. <http://dx.doi.org/10.1212/WNL.000000000000207292>.
- [17] Raffaelli B, Terhart M, Overeem LH, Mecklenburg J, Neeb L, Steinicke M, et al. Migraine evolution after the cessation of CGRP(-receptor) antibody prophylaxis: a prospective, longitudinal cohort study. *Cephalalgia Int J Headache* 2022;42:326–34. <http://dx.doi.org/10.1177/03331024211046617>.
- [18] Reuter U, Ehrlich M, Gendolla A, Heinze A, Klatt J, Wen S, et al. Erenumab versus topiramate for the prevention of migraine – a randomised, double-blind, active-controlled phase 4 trial. *Cephalalgia Int J Headache* 2022;42:108–18. <http://dx.doi.org/10.1177/03331024211053571>.

- [19] Frank F, Ulmer H, Sidoroff V, Broessner G. CGRP-antibodies, topiramate and botulinum toxin type A in episodic and chronic migraine: a systematic review and meta-analysis. *Cephalalgia Int J Headache* 2021;41:1222–39. <http://dx.doi.org/10.1177/03331024211018137>.
- [20] Pozo-Rosich P, Dolezil D, Paemeleire K, Stepien A, Stude P, Snellman J, et al. Early use of erenumab vs. nonspecific oral migraine preventives: the APPRAISE Randomized Clinical Trial. *JAMA Neurol* 2024;81:461–70. <http://dx.doi.org/10.1001/jamaneurol.2024.0368>.
- [21] Hong JB, Lange KS, Overeem LH, Triller P, Raffaelli B, Reuter U. A scoping review and meta-analysis of anti-CGRP monoclonal antibodies: predicting response. *Pharm Basel Switz* 2023;16:934. <http://dx.doi.org/10.3390/ph16070934>.
- [22] Caronna E, Gallardo VJ, Egeo G, Vázquez MM, Castellanos CN, Membrilla JA, et al. Redefining migraine prevention: early treatment with anti-CGRP monoclonal antibodies enhances response in the real world. *J Neurol Neurosurg Psychiatry* 2024;81:461–70. <http://dx.doi.org/10.1136/jnnp-2023-333295>.
- [23] Lanteri-Minet M, Fabre R, Martin C, Pradat K, Alchaar A, Bozzolo E, et al. One-year prospective real-world assessment of effectiveness and safety of erenumab in migraine prevention: results of the French FHU INOVAIN registry study. *J Headache Pain* 2023;24:152. <http://dx.doi.org/10.1186/s10194-023-01680-4>.
- [24] Lee MJ, Al-Karagholi MA-M, Reuter U. New migraine prophylactic drugs: current evidence and practical suggestions for non-responders to prior therapy. *Cephalalgia Int J Headache* 2023;43. <http://dx.doi.org/10.1177/03331024221146315> [3331024221146315].
- [25] Mechtler L, Saikali N, McVige J, Hughes O, Traut A, Adams AM. Real-world evidence for the safety and efficacy of CGRP monoclonal antibody therapy added to OnabotulinumtoxinA treatment for migraine prevention in adult patients with chronic migraine. *Front Neurol* 2021;12:788159. <http://dx.doi.org/10.3389/fneur.2021.788159>.
- [26] Al-Hassany L, Lyons HS, Boucherie DM, Farham F, Lange KS, Marschollek K, et al. The sense of stopping migraine prophylaxis. *J Headache Pain* 2023;24:9. <http://dx.doi.org/10.1186/s10194-023-01539-8>.
- [27] Iannone LF, Fattori D, Benemei S, Chiarugi A, Geppetti P, De Cesaris F. Predictors of sustained response and effects of the discontinuation of anti-calcitonin gene related peptide antibodies and reinitiation in resistant chronic migraine. *Eur J Neurol* 2022;29:1505–13. <http://dx.doi.org/10.1111/ene.15260>.
- [28] Vernieri F, Brunelli N, Messina R, Costa CM, Colombo B, Torelli P, et al. Discontinuing monoclonal antibodies targeting CGRP pathway after one-year treatment: an observational longitudinal cohort study. *J Headache Pain* 2021;22:154. <http://dx.doi.org/10.1186/s10194-021-01363-y>.
- [29] Vernieri F, Brunelli N, Guerzoni S, Iannone LF, Baraldi C, Rao R, et al. Retreating migraine patients in the second year with monoclonal antibodies anti-CGRP pathway: the multicenter prospective cohort RE-DO study. *J Neurol* 2023. <http://dx.doi.org/10.1007/s00415-023-11872-2>.
- [30] de Vries Lentsch S, van der Arend BWH, Maassen VanDenBrink A, Terwindt GM. Blood pressure in patients with migraine treated with monoclonal anti-CGRP (receptor) antibodies: a prospective follow-up study. *Neurology* 2022;99:e1897–904. <http://dx.doi.org/10.1212/WNL.0000000000201008>.
- [31] Reuter U, Goadsby PJ, Ferrari MD, Da Silva Lima GP, Mondal S, Kalim J, et al. Efficacy and safety of erenumab in participants with episodic migraine in whom 2-4 prior preventive treatments had failed: LIBERTY 3-Year Study. *Neurology* 2024;102:e209349. <http://dx.doi.org/10.1212/WNL.0000000000209349>.
- [32] Gérard AO, Merino D, Van Obberghen EK, Rocher F, Destere A, Lanteri-Minet M, et al. Calcitonin gene-related peptide-targeting drugs and Raynaud's phenomenon: a real-world potential safety signal from the WHO pharmacovigilance database. *J Headache Pain* 2022;23:53. <http://dx.doi.org/10.1186/s10194-022-01424-w>.
- [33] Ruiz M, Cocores A, Tosti A, Goadsby PJ, Monteith TS. Alopecia as an emerging adverse event to CGRP monoclonal antibodies: cases series, evaluation of FAERS, and literature review. *Cephalalgia Int J Headache* 2023;43. <http://dx.doi.org/10.1177/03331024221143538> [3331024221143538].
- [34] Al-Hassany L, Vries T de, Carpay JA, MaassenVanDenBrink A. Could erectile dysfunction be a side effect of CGRP inhibition? A case report. *Cephalalgia Int J Headache* 2022;42:257–61. <http://dx.doi.org/10.1177/03331024211037304>.
- [35] Al-Hassany L, Boucherie DM, Couturier EGM, MaassenVanDenBrink A. Case reports: could sexual dysfunction in women with migraine be a side effect of CGRP inhibition? *Cephalalgia Int J Headache* 2024;44. <http://dx.doi.org/10.1177/03331024241248837> [3331024241248837].
- [36] Wurthmann S, Nägel S, Hadaschik E, Schlott S, Scheffler A, Kleinschnitz C, et al. Impaired wound healing in a migraine patient as a possible side effect of calcitonin gene-related peptide receptor antibody treatment: a case report. *Cephalalgia Int J Headache* 2020;40:1255–60. <http://dx.doi.org/10.1177/0333102420933571>.
- [37] Nosedà R, Bedussi F, Gobbi C, Ceschi A, Zecca C. Safety profile of monoclonal antibodies targeting the calcitonin gene-related peptide system in pregnancy: updated analysis in VigiBase®. *Cephalalgia Int J Headache* 2023;43. <http://dx.doi.org/10.1177/03331024231158083> [3331024231158083].
- [38] Sun W, Li Y, Xia B, Chen J, Liu Y, Pang J, et al. Adverse event reporting of four anti-calcitonin gene-related peptide monoclonal antibodies for migraine prevention: a real-world study based on the FDA adverse event reporting system. *Front Pharmacol* 2024;14:1257282. <http://dx.doi.org/10.3389/fphar.2023.1257282>.
- [39] Reuter U, Goadsby PJ, Lanteri-Minet M, Wen S, Hours-Zesiger P, Ferrari MD, et al. Efficacy and tolerability of erenumab in patients with episodic migraine in whom two-to-four previous preventive treatments were unsuccessful: a randomised, double-blind, placebo-controlled, phase 3b study. *Lancet Lond Engl* 2018;392:2280–7. [http://dx.doi.org/10.1016/S0140-6736\(18\)32534-0](http://dx.doi.org/10.1016/S0140-6736(18)32534-0).
- [40] Ferrari MD, Diener HC, Ning X, Galic M, Cohen JM, Yang R, et al. Fremanezumab versus placebo for migraine prevention in patients with documented failure to up to four migraine preventive medication classes (FOCUS): a randomised, double-blind, placebo-controlled, phase 3b trial. *Lancet Lond Engl* 2019;394:1030–40. [http://dx.doi.org/10.1016/S0140-6736\(19\)31946-4](http://dx.doi.org/10.1016/S0140-6736(19)31946-4).
- [41] Mulleners WM, Kim B-K, Láinez MJA, Lanteri-Minet M, Pozo-Rosich P, Wang S, et al. Safety and efficacy of galcanezumab in patients for whom previous migraine preventive medication from two to four categories had failed (CONQUER): a multicentre, randomised, double-blind, placebo-controlled, phase 3b trial. *Lancet Neurol* 2020;19:814–25. [http://dx.doi.org/10.1016/S1474-4422\(20\)30279-9](http://dx.doi.org/10.1016/S1474-4422(20)30279-9).
- [42] Ashina M, Lanteri-Minet M, Pozo-Rosich P, Ettrup A, Christoffersen CL, Josiassen MK, et al. Safety and efficacy of eptinezumab for migraine prevention in patients with two-to-four previous preventive treatment failures (DELIVER): a multi-arm, randomised, double-blind, placebo-controlled, phase 3b trial. *Lancet Neurol* 2022;21:597–607. [http://dx.doi.org/10.1016/S1474-4422\(22\)00185-5](http://dx.doi.org/10.1016/S1474-4422(22)00185-5).

- [43] Lipton RB, Croop R, Stock EG, Stock DA, Morris BA, Frost M, et al. Rimegepant, an oral calcitonin gene-related peptide receptor antagonist, for migraine. *N Engl J Med* 2019;381:142–9. <http://dx.doi.org/10.1056/NEJMoa1811090>.
- [44] Croop R, Lipton RB, Kudrow D, Stock DA, Kamen L, Conway CM, et al. Oral rimegepant for preventive treatment of migraine: a phase 2/3, randomised, double-blind, placebo-controlled trial. *Lancet Lond Engl* 2021;397:51–60. [http://dx.doi.org/10.1016/S0140-6736\(20\)32544-7](http://dx.doi.org/10.1016/S0140-6736(20)32544-7).
- [45] Ailani J, Lipton RB, Goadsby PJ, Guo H, Miceli R, Severt L, et al. Atogepant for the preventive treatment of migraine. *N Engl J Med* 2021;385:695–706. <http://dx.doi.org/10.1056/NEJMoa2035908>.
- [46] Pozo-Rosich P, Ailani J, Ashina M, Goadsby PJ, Lipton RB, Reuter U, et al. Atogepant for the preventive treatment of chronic migraine (PROGRESS): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Lond Engl* 2023;402:775–85. [http://dx.doi.org/10.1016/S0140-6736\(23\)01049-8](http://dx.doi.org/10.1016/S0140-6736(23)01049-8).
- [47] Tassorelli C, Nagy K, Pozo-Rosich P, Lanteri-Minet M, Sacco S, Nežádal T, et al. Safety and efficacy of atogepant for the preventive treatment of episodic migraine in adults for whom conventional oral preventive treatments have failed (ELEVATE): a randomised, placebo-controlled, phase 3b trial. *Lancet Neurol* 2024;23:382–92. [http://dx.doi.org/10.1016/S1474-4422\(24\)00025-5](http://dx.doi.org/10.1016/S1474-4422(24)00025-5).
- [48] Lipton RB, Croop R, Stock DA, Madonia J, Forshaw M, Lovegren M, et al. Safety, tolerability, and efficacy of zavegepant 10 mg nasal spray for the acute treatment of migraine in the USA: a phase 3, double-blind, randomised, placebo-controlled multicentre trial. *Lancet Neurol* 2023;22:209–17. [http://dx.doi.org/10.1016/S1474-4422\(22\)00517-8](http://dx.doi.org/10.1016/S1474-4422(22)00517-8).
- [49] Dodick DW, Lipton RB, Ailani J, Lu K, Finnegan M, Trugman JM, et al. Ubrogepant for the treatment of migraine. *N Engl J Med* 2019;381:2230–41. <http://dx.doi.org/10.1056/NEJMoa1813049>.
- [50] Dodick DW, Goadsby PJ, Schwedt TJ, Lipton RB, Liu C, Lu K, et al. Ubrogepant for the treatment of migraine attacks during the prodrome: a phase 3, multicentre, randomised, double-blind, placebo-controlled, crossover trial in the USA. *Lancet Lond Engl* 2023;402:2307–16. [http://dx.doi.org/10.1016/S0140-6736\(23\)01683-5](http://dx.doi.org/10.1016/S0140-6736(23)01683-5).
- [51] Saengjaroenatham C, Strother LC, Dripps I, Sultan Jabir MR, Pradhan A, Goadsby PJ, et al. Differential medication overuse risk of novel anti-migraine therapeutics. *Brain J Neurol* 2020;143:2681–8. <http://dx.doi.org/10.1093/brain/awaa211>.
- [52] Navratilova E, Behraves S, Oyarzo J, Dodick DW, Banerjee P, Porreca F. Ubrogepant does not induce latent sensitization in a preclinical model of medication overuse headache. *Cephalalgia Int J Headache* 2020;40:892–902. <http://dx.doi.org/10.1177/0333102420938652>.
- [53] Sacco S, Lampl C, Amin FM, Braschinsky M, Deligianni C, Uludüz D, et al. European Headache Federation (EHF) consensus on the definition of effective treatment of a migraine attack and of triptan failure. *J Headache Pain* 2022;23:133. <http://dx.doi.org/10.1186/s10194-022-01502-z>.
- [54] Ruscheweyh R, Gossrau G, Dresler T, Freilinger T, Förderreuther S, Gaul C, et al. Triptan non-response in specialized headache care: cross-sectional data from the DMKG Headache Registry. *J Headache Pain* 2023;24:135. <http://dx.doi.org/10.1186/s10194-023-01676-0>.
- [55] Law S, Derry S, Moore RA. Sumatriptan plus naproxen for the treatment of acute migraine attacks in adults. *Cochrane Database Syst Rev* 2016;4:CD008541. <http://dx.doi.org/10.1002/14651858.CD008541.pub3>.
- [56] Lipton RB, Blumenfeld A, Jensen CM, Croop R, Thiry A, L'Italien G, et al. Efficacy of rimegepant for the acute treatment of migraine based on triptan treatment experience: Pooled results from three phase 3 randomized clinical trials. *Cephalalgia Int J Headache* 2023;43. <http://dx.doi.org/10.1177/03331024221141686>. [3331024221141686].
- [57] Naghdi S, Underwood M, Madan J, Brown A, Duncan C, Matharu M, et al. Clinical effectiveness of pharmacological interventions for managing chronic migraine in adults: a systematic review and network meta-analysis. *J Headache Pain* 2023;24:164. <http://dx.doi.org/10.1186/s10194-023-01696-w>.
- [58] Lanteri-Minet M, Ducros A, Francois C, Olewinska E, Nikodem M, Dupont-Benjamin L. Effectiveness of onabotulinumtoxinA (BOTOX®) for the preventive treatment of chronic migraine: a meta-analysis on 10 years of real-world data. *Cephalalgia Int J Headache* 2022;42:1543–64. <http://dx.doi.org/10.1177/03331024221123058>.
- [59] Brin MF, Kirby RS, Slavotinek A, Adams AM, Parker L, Ukah A, et al. Pregnancy outcomes in patients exposed to onabotulinumtoxinA treatment: a cumulative 29-year safety update. *Neurology* 2023;101:e103–13. <http://dx.doi.org/10.1212/WNL.0000000000207375>.
- [60] Bjørk M-H, Zoega H, Leinonen MK, Cohen JM, Dreier JW, Furu K, et al. Association of prenatal exposure to antiepileptic medication with risk of autism and intellectual disability. *JAMA Neurol* 2022;79:672–81. <http://dx.doi.org/10.1001/jamaneurol.2022.1269>.
- [61] Actualité – Topiramate et risques chez les enfants exposés pendant la grossesse : modification des conditions de prescription et de délivrance aux femmes concernées. ANSM n.d. <https://ansm.sante.fr/actualites/topiramate-et-risques-chez-les-enfants-exposes-pendant-la-grossesse-modification-des-conditions-de-prescription-et-de-delivrance-aux-femmes-concernees-5> [accessed February 15, 2024].
- [62] Hernández-Díaz S, Straub L, Bateman BT, Zhu Y, Mogun H, Wisner KL, et al. Risk of autism after prenatal topiramate, valproate, or lamotrigine exposure. *N Engl J Med* 2024;390:1069–79. <http://dx.doi.org/10.1056/NEJMoa2309359>.
- [63] Dreier JW, Bjørk M-H, Alvestad S, Gissler M, Igland J, Leinonen MK, et al. Prenatal exposure to antiepileptic medication and incidence of childhood- and adolescence-onset psychiatric disorders. *JAMA Neurol* 2023;80:568–77. <http://dx.doi.org/10.1001/jamaneurol.2023.0674>.
- [64] Chowdhury D, Bansal L, Duggal A, Datta D, Mundra A, Krishnan A, et al. TOP-PRO study: a randomized double-blind controlled trial of topiramate versus propranolol for prevention of chronic migraine. *Cephalalgia Int J Headache* 2022;42:396–408. <http://dx.doi.org/10.1177/03331024211047454>.
- [65] Information de sécurité – Valproate et dérivés: risque potenti. ANSM n.d. <https://ansm.sante.fr/informations-de-securite/valproate-et-derives-risque-potentiel-de-troubles-neurodeveloppementaux-chez-les-enfants-dont-le-pere-a-ete-traite-dans-les-3-mois-qui-precedent-la-conception> [accessed October 16, 2023].
- [66] Christensen J, Trabjerg BB, Dreier JW. Valproate use during spermatogenesis and risk to offspring. *JAMA Netw Open* 2024;7:e2414709. <http://dx.doi.org/10.1001/jamanetworkopen.2024.14709>.
- [67] Deligianni CI, Sacco S, Ekizoglu E, Uluduz D, Gil-Gouveia R, MaassenVanDenBrink A, et al. European Headache Federation (EHF) critical re-appraisal and meta-analysis of oral drugs in migraine prevention-part 2: flunarizine. *J Headache Pain* 2023;24:128. <http://dx.doi.org/10.1186/s10194-023-01657-3>.